



Micro: OFC 20-25 cm (8-10 in

Instructions for use:

- 1. Disconnect the Velcro.
- 2. Position the eye pads over the eyes.
- 3. Pass the lower strap under the small of the neck and attach velcro.
- Secure the Velcro of both straps and position the EyeMax2™ as shown.
- Make sure the strap completely covers the ears.
- 6. If the eye patch rides up away from the nose, tighten the lower strap.

Catalog # R300P01 Regular: occipital-frontal circumference 33-38 cm in packs of 20. Catalog # R300P02 Preemie: occipital-frontal circumference 26-32 cm in packs of 20. Catalog # R300P03 Micro: occipital-frontal circumference 20-25 cm in packs of 20.



Single Patient Use Only Non-Sterile, Latex-Free



Precautions:

Caution: Do not tug on the EyeMax™ to reposition it.

Caution: Do not tug on the EyeMax[™] to reposition it.

A premature infant's skin is very delicate and can easily rupture.

Caution: Avoid obstructing the nose. Check regularly to ensure the EyeMax[™] is not blocking the patient's airway.

Caution: The EyeMax[™] is not a bandage. Cover any existing skin abrasions

with appropriate bandages before use.

Caution: Do not over tighten – causing excessive pressure on the eyes.

Caution: Make sure the strap is covering the ear, not hooked over the ear, and

that the ear is not folded over.

Caution: Replace the EyeMax™ with a new one if it becomes over-stretched.

Preemie: OFC 26-32 cm (10-13 in.)

Instructions for use:

- 1. Disconnect the Velcro.
- 2. Position the eye pads over the eyes.
- 3. Pass the lower strap under the small of the neck and attach velcro.
- Secure the Velcro of both straps and position the EyeMax2™ as shown.
- 5. Make sure the strap completely covers the ears.
- 6. If the eye patch rides up away from the nose, tighten the lower strap.

Catalog # R300P01 Regular: occipital-frontal circumference 33-38 cm in packs of 20. Catalog # R300P02 Preemie: occipital-frontal circumference 26-32 cm in packs of 20. Catalog # R300P03 Micro: occipital-frontal circumference 20-25 cm in packs of 20.



Single Patient Use Only Non-Sterile, Latex-Free



Regular: OFC 33-38 cm (13-15 in.)

Instructions for use:

- 1. Disconnect the Velcro.
- 2. Position the eye pads over the eyes
- 3. Pass the lower strap under the small of the neck and attach velcro. Secure the Velcro of both straps and position the EyeMax2™ as shown.
- 5. Make sure the strap completely covers the ears.
- 6. If the eye patch rides up away from the nose, tighten the lower strap.

Catalog # R300P01 **Regular**: occipital-frontal circumference **33-38** cm in packs of 20. Catalog # R300P02 **Preemie**: occipital-frontal circumference **26-32** cm in packs of 20. Catalog # R300P03 Micro: occipital-frontal circumference 20-25 cm in packs of 20.



Single Patient Use Only Non-Sterile, Latex-Free



6526 South Cottonwood Street Salt Lake City, Utah 84107

Precautions:

Caution: Do not tug on the EyeMax™ to reposition it.

A premature infant's skin is very delicate and can easily rupture.

Caution: Avoid obstructing the nose. Check regularly to ensure the

EyeMax™ is not blocking the patient's airway.

Caution: The EyeMax™ is not a bandage. Cover any existing skin abrasions with appropriate bandages before use.

Caution: Do not over tighten - causing excessive pressure on the eyes.

Caution: Make sure the strap is covering the ear, not hooked over the ear, and that the ear is not folded over.

Caution: Replace the EyeMax™ with a new one if it becomes over-stretched.

Precautions:



Caution: Do not tug on the EyeMax™ to reposition it.

A premature infant's skin is very delicate and can easily rupture. Caution: Avoid obstructing the nose. Check regularly to ensure the

EyeMax™ is not blocking the patient's airway.

Caution: The EyeMax™ is not a bandage. Cover any existing skin abrasions

with appropriate bandages before use.

Caution: Do not over tighten – causing excessive pressure on the eyes.

Caution: Make sure the strap is covering the ear, not hooked over the ear, and that the ear is not folded over

Caution: Replace the EyeMax™ with a new one if it becomes over-stretched.

Distributed in the UK by:



Viamed Limited · 15 Station Road · Cross Hills Keighley · West Yorkshire · BD20 7DT · United Kingdom Tel: +44 (0)1535 634 542 Fax: +44 (0)1535 635 582 Email: info@viamed.co.uk Website: www.viamed.co.uk